

On prime factors of class number of cyclotomic fields

Roland Quême

2006 jan 09

Contents

1	Some definitions	2
2	On Kummer and Stickelberger relation	3
3	Polynomial congruences mod p connected to the p-class group C_p	6
3.1	Stickelberger relation for prime ideals $\mathfrak{q}$ of inertial degree $f = 1$	6
3.2	Stickelberger relation for prime ideals $\mathfrak{q}$ of inertial degree $f > 1$	7
3.3	Polynomial congruences mod p^2 connected to the p^2 -class group C_{p^2}	9
4	On prime factors $h \neq p$ of the class number of the p-cyclotomic field	11
4.1	The general case	12
4.2	The case of complex quadratic fields contained in K	15
4.3	The case of biquadratic fields contained in K	17
5	A numerical MAPLE algorithm	18

Abstract

Let p be an odd prime. Let $K = \mathbb{Q}(\zeta)$ be the p -cyclotomic number field. Let v be a primitive root mod p and $\sigma : \zeta \rightarrow \zeta^v$ be a $\mathbb{Q}$ -isomorphism of the extension $K/\mathbb{Q}$ generating the Galois group G of $K/\mathbb{Q}$. Following the conventions of Ribenboim in [8], for $n \in \mathbb{Z}$, the notation v_n is understood by $v_n = v^n \bmod p$ with $1 \leq v_n \leq p - 1$. Let $P(X) = \sum_{i=0}^{p-2} X^i v_{-i} \in \mathbb{Z}[X]$ be the Stickelberger polynomial. $P(\sigma)$ annihilates the class group $\mathbf{C}$ of K . There exists a polynomial $Q(X) \in \mathbb{Z}[G]$ such that $P(\sigma) \times (\sigma - v) = p \times Q(\sigma)$ and such that $Q(\sigma)$ annihilates the p -class group C_p of K .

These result allow:

1. to describe the structure of the relative class group $\mathbf{C}^-$,
2. to give some explicit congruences in $\mathbb{Z}[v] \bmod p$ for the p -class group of K (the subgroup of exponent p of $\mathbf{C}$),
3. to give some explicit congruences in $\mathbb{Z}[v] \bmod h$ for the h -class group of K for the prime divisors $h \neq p$ of the class number $h(K)$.
4. We detail at the end the case of class number of quadratic and biquadratic fields contained in the cyclotomic field K .
5. As application, we give a MAPLE algorithm which describes the structure of the relative class group $\mathbf{C}^-$ of the cyclotomic field K for all the prime numbers $p < 500$.

This article is at elementary level.

Remark: we have not found in the literature some formulations corresponding to theorems 3.2 p. 7, 4.1 p. 12, 4.3 p. 13, 4.6 p. 16, 4.7 p. 17, and 4.8 p. 17 and to MAPLE algorithm described in section 5 p 18.

1 Some definitions

In this section we give some definitions and notations on cyclotomic fields and p -class group used in this paper.

1. Let p be an odd prime. Let $\mathbf{F}_p$ be the finite field of cardinal p and $\mathbf{F}_p^*$ its multiplicative group. Let ζ be a root of the polynomial equation $X^{p-1} + X^{p-2} + \dots + X + 1 = 0$. Let K be the p -cyclotomic field $K = \mathbb{Q}(\zeta)$ and O_K its ring of integers. Let K^+ be the maximal totally real subfield of K . Let v be a primitive root $\bmod p$. In this paper, following Ribenboim conventions in Ribenboim [8] for any $n \in \mathbb{Z}$, we note $v_n = v^n \bmod p$ with $1 \leq v_n \leq p-1$. Let G be the Galois group of the extension $K/\mathbb{Q}$. Let $\sigma : \zeta \rightarrow \zeta^v$ be a $\mathbb{Q}$ -isomorphism of the extension $K/\mathbb{Q}$ generating G . Let $\lambda = \zeta - 1$. The prime ideal of K lying over p is $\pi = \lambda O_K$.
2. Let $\mathbf{C}$ be the class group of K . Let C_p be the p -class of K (the subgroup of exponent p of $\mathbf{C}$). Let C_p^+ be the p -class group of K^+ . Then $C_p = C_p^+ \oplus C_p^-$ where C_p^- is called the relative p -class group. Let r^- be the rank of C_p^- .
3. C_p^- is the direct sum of r^- subgroups Γ_k of order p annihilated by $\sigma - \mu_k \in \mathbf{F}_p[G]$ with $\mu_k \in \mathbf{F}_p^*$, $\mu_k = v_{2m_k+1}$ where m_k is a natural integer m_k , $1 \leq m_k \leq \frac{p-3}{2}$,

$$(1) \quad C_p^- = \bigoplus_{k=1}^{r^-} \Gamma_k.$$

2 On Kummer and Stickelberger relation

Stickelberger relation was already known by Kummer under the form of Jacobi resolvents for the cyclotomic field K , see for instance Ribenboim [8] (2.6) p. 119. In this section we derive some elementary properties from Stickelberger relation.

1. Let $q \neq p$ be an odd prime. Let ζ_q be a root of the minimal polynomial equation $X^{q-1} + X^{q-2} + \dots + X + 1 = 0$. Let $K_q = \mathbb{Q}(\zeta_q)$ be the q -cyclotomic field. Let $K_{pq} = \mathbb{Q}(\zeta_p, \zeta_q)$. Then K_{pq} is the compositum KK_q . The ring of integers of K_{pq} is $O_{K_{pq}}$.
2. Let $\mathfrak{q}$ be a prime ideal of O_K lying over the prime q . Let f be the order of $q \bmod p$ and $m = N_{K_p/\mathbb{Q}}(\mathfrak{q}) = q^f$. If $\psi(\alpha) = a$ is the image of $\alpha \in O_K$ under the natural map $\psi : O_K \rightarrow O_K/\mathfrak{q}$, then for $\psi(\alpha) = a \not\equiv 0$ define a character $\chi_{\mathfrak{q}}^{(p)}$ on $\mathbf{F}_m = O_K/\mathfrak{q}$ by

$$(2) \quad \chi_{\mathfrak{q}}^{(p)}(a) = \left\{ \frac{a}{\mathfrak{q}} \right\}_p^{-1} = \overline{\left\{ \frac{a}{\mathfrak{q}} \right\}_p},$$

where $\left\{ \frac{a}{\mathfrak{q}} \right\} = \zeta^c$ for some natural integer c , is the p^{th} power residue character mod $\mathfrak{q}$. We define the Gauss sum

$$(3) \quad g(\mathfrak{q}) = \sum_{x \in \mathbf{F}_m} (\chi_{\mathfrak{q}}^{(p)}(x) \times \zeta_q^{Tr_{\mathbf{F}_m/\mathbf{F}_q}(x)}) \in O_{K_{pq}}.$$

It follows that $g(\mathfrak{q}) \in O_{K_{pq}}$. Moreover $g(\mathfrak{q})^p \in O_K$, see for instance Mollin [6] prop. 5.88 (c) p. 308.

3. The Stickelberger relation is classically:

$$(4) \quad g(\mathfrak{q})^p O_K = \mathfrak{q}^S,$$

with $S = \sum_{t=1}^{p-1} t \times \varpi_t^{-1}$, where $\varpi_t \in Gal(K/\mathbb{Q})$ is given by $\varpi_t : \zeta \rightarrow \zeta^t$ (see for instance Mollin [6] thm. 5.109 p. 315).

The four following lemmas are derived in an elementary way from the Stickelberger relation.

Lemma 2.1. *If $q \not\equiv 1 \bmod p$ then the Gauss sum $g(\mathfrak{q}) \in \mathbb{Z}[\zeta]$.*

Proof.

1. Let u be a primitive root mod q . Let $\tau : \zeta_q \rightarrow \zeta_q^u$ be a $\mathbb{Q}$ -isomorphism generating $Gal(K_q/\mathbb{Q})$. The $\mathbb{Q}$ -isomorphism τ is extended to a K_p -isomorphism of K_{pq} by $\tau : \zeta_q \rightarrow \zeta_q^u, \quad \zeta_p \rightarrow \zeta_p$. Then $g(\mathfrak{q})^p \in \mathbb{Z}[\zeta_p]$ and so

$$\tau(g(\mathfrak{q}))^p = g(\mathfrak{q})^p,$$

and it follows that there exists a natural integer ρ with $\rho < p$ such that

$$\tau(g(\mathbf{q})) = \zeta_p^\rho \times g(\mathbf{q}).$$

Then $N_{K_{pq}/K}(\tau(g(\mathbf{q}))) = \zeta^{(q-1)\rho} \times N_{K_{pq}/K}(g(\mathbf{q}))$ and so $\zeta^{\rho(q-1)} = 1$.

2. If $q \not\equiv 1 \pmod{p}$, it implies that $\zeta^\rho = 1$ and so that $\tau(g(\mathbf{q})) = g(\mathbf{q})$ and thus that $g(\mathbf{q}) \in O_K$.

□

Lemma 2.2. *Let $S = \sum_{t=1}^{p-1} \varpi_t^{-1} \times t$ where ϖ_t is the $\mathbb{Q}$ -isomorphism of the extension $K/\mathbb{Q}$ given by $\varpi_t : \zeta \rightarrow \zeta^t$ of K . Let $P(\sigma) = \sum_{i=0}^{p-2} \sigma^i \times v_{-i} \in \mathbb{Z}[G]$. Then $S = P(\sigma)$.*

Proof. Let us consider one term $\varpi_t^{-1} \times t$. Then $v_{-1} = v_{p-2}$ is a primitive root mod p and so there exists one and one i such that $t = v_{-i}$. Then $\varpi_{v_{-i}} : \zeta \rightarrow \zeta^{v_{-i}}$ and so $\varpi_{v_{-i}}^{-1} : \zeta \rightarrow \zeta^{v_i}$ and so $\varpi_{v_{-i}}^{-1} = \sigma^i$ (observe that $\sigma^{p-1} \times v_{-(p-1)} = 1$), which achieves the proof. □

Lemma 2.3.

$$(5) \quad P(\sigma) = \sum_{i=0}^{p-2} \sigma^i \times v_{-i} = v_{-(p-2)} \times \left\{ \prod_{k=0, k \neq 1}^{p-2} (\sigma - v_k) \right\} + p \times R(\sigma),$$

where $R(\sigma) \in \mathbb{Z}[G]$ with $\deg(R(\sigma)) < p - 2$.

Proof. Let us consider the polynomial $R_0(\sigma) = P(\sigma) - v_{-(p-2)} \times \left\{ \prod_{k=0, k \neq 1}^{p-2} (\sigma - v_k) \right\}$ in $\mathbf{F}_p[G]$. Then $R_0(\sigma)$ is of degree smaller than $p - 2$ and the two polynomials $\sum_{i=0}^{p-2} \sigma^i v_{-i}$ and $\prod_{k=0, k \neq 1}^{p-2} (\sigma - v_k)$ take a null value in $\mathbf{F}_p[G]$ when σ takes the $p - 2$ different values $\sigma = v_k$ for $k = 0, \dots, p - 2$, $k \neq 1$. Then $R_0(\sigma) = 0$ in $\mathbf{F}_p[G]$ which leads to the result in $\mathbb{Z}[G]$. □

Lemma 2.4.

$$(6) \quad P(\sigma) \times (\sigma - v) = p \times Q(\sigma),$$

where $Q(\sigma) = \sum_{i=1}^{p-2} \delta_i \times \sigma^i \in \mathbb{Z}[G]$ is given by

$$\begin{aligned}
 \delta_{p-2} &= \frac{v_{-(p-3)} - v_{-(p-2)}v}{p}, \\
 \delta_{p-3} &= \frac{v_{-(p-4)} - v_{-(p-3)}v}{p}, \\
 &\vdots \\
 \delta_i &= \frac{v_{-(i-1)} - v_{-i}v}{p}, \\
 &\vdots \\
 \delta_1 &= \frac{1 - v_{-1}v}{p},
 \end{aligned}
 \tag{7}$$

with $-p < \delta_i \leq 0$.

Proof. We start of the relation in $\mathbb{Z}[G]$

$$P(\sigma) \times (\sigma - v) = v_{-(p-2)} \times \prod_{k=0}^{p-2} (\sigma - v_k) + p \times R(\sigma) \times (\sigma - v) = p \times Q(\sigma),$$

with $Q(\sigma) \in \mathbb{Z}[G]$ because $\prod_{k=0}^{p-2} (\sigma - v_k) = 0$ in $\mathbf{F}_p[G]$ and so $\prod_{k=0}^{p-2} (\sigma - v_k) = p \times R_1(\sigma)$ in $\mathbb{Z}[G]$. Then we identify in $\mathbb{Z}[G]$ the coefficients in the relation

$$\begin{aligned}
 &(v_{-(p-2)}\sigma^{p-2} + v_{-(p-3)}\sigma^{p-3} + \cdots + v_{-1}\sigma + 1) \times (\sigma - v) = \\
 &p \times (\delta_{p-2}\sigma^{p-2} + \delta_{p-3}\sigma^{p-3} + \cdots + \delta_1\sigma + \delta_0),
 \end{aligned}$$

where $\sigma^{p-1} = 1$. □

Remarks:

1. Observe that we have more generally for the indeterminate X the algebraic identity in $\mathbb{Z}[X]$

$$(8) \quad P(X)(X - v) = p \times Q(X) + v(X^{p-1} - 1).$$

2. Observe that, with our notations, $\delta_i \in \mathbb{Z}$, $i = 1, \dots, p-2$, but generally $\delta_i \not\equiv 0 \pmod{p}$.
3. We see also that $-p < \delta_i \leq 0$. Observe also that $\delta_0 = \frac{v_{-(p-2)} - v}{p} = 0$.

3 Polynomial congruences mod p connected to the p -class group C_p

We give some explicit polynomial congruences in $\mathbb{Z}[v]$ mod p connected to the relative p -class group C_p^- of O_K . We apply successively the Stickelberger relation to prime ideals $\mathfrak{q}$ of inertial degree $f = 1$ and of inertial degree $f > 1$. We recall that r^- is the p -rank of the relative p -class group C_p^- of K .

3.1 Stickelberger relation for prime ideals $\mathfrak{q}$ of inertial degree $f = 1$

Theorem 3.1. $\prod_{k=1}^{r^-} (\sigma - v_{2m_k+1})$ divides $Q(\sigma)$ in $\mathbf{F}_p[G]$ and

$$(9) \quad Q(v_{2m_k+1}) = \sum_{i=1}^{p-2} v_{(2m_k+1) \times i} \times \left(\frac{v_{-(i-1)} - v_{-i} \times v}{p} \right) \equiv 0 \pmod{p},$$

where m_k is defined in relation (1) p. 2.

Proof. From Kummer, the group of ideal classes of K is generated by the classes of prime ideals of degree 1 (see for instance Ribenboim [8] (3A) p. 119). Let $\mathfrak{q}$ a prime ideal of inertial degree 1 whose class $Cl(\mathfrak{q}) \in C_p^-$ is annihilated by $\sigma - \mu_k$ with $\mu_k = v_{2m_k+1}$. We start of $g(\mathfrak{q})^p O_K = \mathfrak{q}^{P(\sigma)}$ and so $g(\mathfrak{q})^{p(\sigma-v)} O_K = \mathfrak{q}^{P(\sigma)(\sigma-v)} = \mathfrak{q}^{pQ(\sigma)}$ and thus $g(\mathfrak{q})^{(\sigma-v)} O_K = \mathfrak{q}^{Q(\sigma)}$. It can be shown that $g(\mathfrak{q})^{\sigma-v} \in K$, see for instance Ribenboim [9] F. p. 440. Therefore $Q(\sigma)$ annihilates the ideal class $Cl(\mathfrak{q})$ and the congruence follows. $\square$

Remarks:

1. Observe that δ_i can also be written in the form $\delta_i = -[\frac{v_{-i} \times v}{p}]$ where $[x]$ is the integer part of x , similar form also known in the literature.
2. Observe that it is possible to get other polynomials of $\mathbb{Z}[G]$ annihilating the relative p -class group C_p^- : for instance from Kummer's formula on Jacobi cyclotomic functions we induce other polynomials $Q_d(\sigma)$ annihilating the relative p -class group C_p^- of K : If $1 \leq d \leq p-2$ define the set

$$I_d = \{i \mid 0 \leq i \leq p-2, \quad v_{(p-1)/2-i} + v_{(p-1)/2-i+ind_v(d)} > p\}$$

where the index $ind_v(d)$ is the minimal integer s such that $d = v_s$. Then the polynomials $Q_d(\sigma) = \sum_{i \in I_d} \sigma^i$ for $d = 1, \dots, p-2$ annihilate the p -class C_p of K , see for instance Ribenboim [8] relations (2.4) and (2.5) p. 119.

3. See also in a more general context Washington, [11] corollary 10.15 p. 198.
4. It is easy to verify the consistency of relation (9) with the table of irregular primes and Bernoulli numbers in Washington, [11] p. 410.
5. See section 5 p. 18 with an algorithm improving this result.

3.2 Stickelberger relation for prime ideals $\mathfrak{q}$ of inertial degree $f > 1$

Let $\mathfrak{q}$ be a prime ideal of O_K with $Cl(\mathfrak{q}) \in C_p$. In this section we apply Stickelberger relation to the prime ideals $\mathfrak{q}$ of inertial degree $f > 1$ with the method used for the prime ideals of inertial degree 1 in section 3 p. 6. Observe, from lemma 2.1 p. 3, that $f > 1$ implies that $g(\mathfrak{q}) \in O_K$, property used in this section (by opposite $g(\mathfrak{q}) \notin K$ when $f = 1$).

A definition: we say that the prime ideal $\mathfrak{c}$ of a number field M is p -principal if the component of the class group $\langle Cl(\mathfrak{c}) \rangle$ in p -class group D_p of M is trivial.

Theorem 3.2. *Let q be an odd prime with $q \neq p$. Let f be the order of $q \bmod p$ and $m = \frac{p-1}{f}$. Let $\mathfrak{q}$ be a prime ideal of O_K lying over q with $Cl(\mathfrak{q}) \in C_p^-$. If $f > 1$ then*

1. $g(\mathfrak{q}) \in O_K$ and $g(\mathfrak{q})O_K = \mathfrak{q}^{P_1(\sigma)}$ where

$$(10) \quad P_1(\sigma) = \sum_{i=0}^{m-1} \left(\frac{\sum_{j=0}^{f-1} v_{-(i+jm)}}{p} \right) \times \sigma^i \in \mathbb{Z}[G].$$

2. *There exists a natural integer n with $1 \leq n \leq p-2$ such that $\sigma - v_n$ divides $P_1(\sigma)$ in $\mathbb{F}_p[G]$ and $\sigma - v_n$ annihilates $Cl(\mathfrak{q})$.*

Proof.

1. $N_{K/\mathbb{Q}}(\mathfrak{q}) = q^f$ and $\mathfrak{q} = \mathfrak{q}^{\sigma^m} = \dots = \mathfrak{q}^{\sigma^{(f-1)m}}$. From Stickelberger relation $g(\mathfrak{q})^p \mathbb{Z}[\zeta_p] = \mathfrak{q}^{P(\sigma)}$ where $P(\sigma) = \sum_{i=0}^{m-1} \sum_{j=0}^{f-1} \sigma^{i+jm} v_{-(i+jm)}$. Observe that, from hypothesis, $\mathfrak{q} = \mathfrak{q}^{\sigma^m} = \dots = \mathfrak{q}^{\sigma^{(f-1)m}}$ so Stickelberger's relation implies that $g(\mathfrak{q})^p O_K = \mathfrak{q}^{P(\sigma)}$ with

$$P(\sigma) = \sum_{i=0}^{m-1} \sum_{j=0}^{f-1} \sigma^i v_{-(i+jm)} = p \times \sum_{i=0}^{m-1} \left(\frac{\sum_{j=0}^{f-1} v_{-(i+jm)}}{p} \right) \times \sigma^i,$$

where $(\sum_{i=0}^{f-1} v_{-(i+jm)})/p \in \mathbb{Z}$ because $v_{-m} - 1 \not\equiv 0 \pmod{p}$.

2. Let $P_1(\sigma) = \frac{P(\sigma)}{p}$. Then from below $P_1(\sigma) \in \mathbb{Z}[G]$. From lemma 2.1 p. 3 we know that $f > 1$ implies that $g(\mathbf{q}) \in O_K$. Therefore

$$g(\mathbf{q})^p O_K = \mathbf{q}^{pP_1(\sigma)}, \quad g(\mathbf{q}) \in O_K,$$

and so

$$g(\mathbf{q}) O_K = \mathbf{q}^{P_1(\sigma)}, \quad g(\mathbf{q}) \in O_K.$$

Then $P_1(\sigma)$ annihilates $Cl(\mathbf{q})$ and thus there exists $1 < n \leq p - 2$ such that $(\sigma - v_n) \mid P_1(\sigma)$.

□

Remarks

1. For $f = 2$ the value of polynomial $P_1(\sigma)$ obtained from this lemma is $P_1(\sigma) = \sum_{i=1}^{(p-3)/2} \sigma^i$.
2. Let $\mathbf{q}$ be a prime not principal ideal of inertial degree $f > 1$ with $Cl(\mathbf{q}) \in C_p$. The **two** polynomials of $\mathbb{Z}[G_p]$, $Q(\sigma) = \sum_{i=0}^{p-2} \left(\frac{v_{-(i-1)} - v_{-i} v}{p} \right) \times \sigma^i$ (see thm 3.1) and $P_1(\sigma) = \sum_{i=0}^{m-1} \left(\frac{\sum_{j=0}^{f-1} v_{-(i+jm)}}{p} \right) \times \sigma^i$ (see lemma 3.2) annihilate the ideal class $Cl(\mathbf{q})$. When $f > 1$ the lemma 3.2 p. 7 supplement the theorem 3.1 p. 6.
3. This result explains that, when f increases, the proportion of p -principal ideals $\mathbf{q}$ increases.

It is possible to derive some explicit congruences in $\mathbb{Z}$ from this theorem.

Corollary 3.3. *Let q be an odd prime with $q \neq p$. Let f be the order of $q \bmod p$ and let $m = \frac{p-1}{f}$. Let $\mathbf{q}$ be an prime ideal of O_K lying over q . Suppose that $f > 1$.*

1. *If the ideal $\mathbf{q}$ is non p -principal there exists a natural integer l , $1 \leq l < m$ such that*

$$(11) \quad \sum_{i=0}^{m-1} \left(\frac{\sum_{j=0}^{f-1} v_{-(i+jm)}}{p} \right) \times v^{lfi} \equiv 0 \bmod p,$$

2. *If for all natural integers l such that $1 \leq l < m$*

$$(12) \quad \sum_{i=0}^{m-1} \left(\frac{\sum_{j=0}^{f-1} v_{-(i+jm)}}{p} \right) \times v^{lfi} \not\equiv 0 \bmod p,$$

then $\mathbf{q}$ is p -principal

Proof.

1. Suppose that $\mathbf{q}$ is not p -principal. Observe at first that congruence (11) with $l = m$ should imply that $\sum_{i=0}^{m-1} (\sum_{j=0}^{f-1} v_{-(i+jm)})/p \equiv 0 \pmod{p}$ or

$$\sum_{i=0}^{m-1} \sum_{j=0}^{f-1} v_{-(i+jm)} \equiv 0 \pmod{p^2},$$

which is not possible because $v_{-(i+jm)} = v_{-(i'+j'm)}$ implies that $j = j'$ and $i = i'$ and so that $\sum_{i=0}^{m-1} \sum_{j=0}^{f-1} v_{-(i+jm)} = \frac{p(p-1)}{2}$.

2. The polynomial $P_1(\sigma)$ of lemma 3.2 annihilates the non p -principal ideal $\mathbf{q}$ in $\mathbf{F}_p[G]$ only if there exists $\sigma - v_n$ dividing $P_1(\sigma)$ in $\mathbf{F}_p[G]$. From $\mathbf{q}^{\sigma^m-1} = 1$ it follows also that $\sigma - v_n \mid \sigma^m - 1$. But $\sigma - v_n \mid \sigma^m - v_{nm}$ and so $\sigma - v_n \mid v_{nm} - 1$, thus $nm \equiv 0 \pmod{p-1}$, so $n \equiv 0 \pmod{f}$ and $n = lf$ for some l . Therefore if $\mathbf{q}$ is non p -principal there exists a natural integer l , $1 \leq l < m$ such that

$$(13) \quad \sum_{i=0}^{m-1} \left(\frac{\sum_{j=0}^{f-1} v_{-(i+jm)}}{p} \right) \times v^{lfi} \equiv 0 \pmod{p},$$

3. The relation (12) is an immediate consequence of previous part of the proof.

□

3.3 Polynomial congruences mod p^2 connected to the p^2 -class group C_{p^2}

Let $C_{p^2}^-$ be the subgroup of exponent p^2 (so with elements of order dividing p^2) of the relative class group $\mathbf{C}^-$ of K .

1. We have seen in relation (1) p. 2 that the relative p -class group C_p^- can be seen as a direct sum $C_p^- = \oplus_{i=1}^{r^-} \Gamma_i$ where Γ_i is a cyclic subgroup of C_p^- annihilated by $\sigma - \mu_i$ with $2 \leq \mu_i \leq p-2$.
2. $C_{p^2}^-$ can be seen as a direct sum

$$(14) \quad C_{p^2}^- = \oplus_{i=1}^{r^-} \Delta_i,$$

where Δ_i is a cyclic group with $\Gamma_i \subset \Delta_i$ and whose order divides p^2 .

3. Suppose that Δ_i is of order p^2 . Show that Δ_i is annihilated by $\sigma - (\mu_i + a_i p)$ with a_i natural integer $1 \leq a_i \leq p-1$:
 - (a) From Kummer, there exist some prime ideals $\mathbf{Q}_i$ of O_K with $Cl(\mathbf{Q}_i) \in \Delta_i$, $Cl(\mathbf{Q}_i^p) \in \Gamma_i$ and $\langle \mathbf{Q}_i^{p(\sigma-\mu_i)} \rangle$ principal as seen in previous sections. Therefore $\langle \mathbf{Q}_i^{\sigma-\mu_i} \rangle$ is of order p .

- (b) $Cl(\mathbf{Q}_i^{\mu_i}) \in \Delta_i$. In the other hand $Cl(\mathbf{Q}_i^\sigma) \in \Delta_i$: if not $Cl(\mathbf{Q}_i^\sigma)$ should have at least one component $\mathbf{c}_j \in \Delta_j$, $j \neq i$ and so $Cl(\mathbf{Q}_i^{p\sigma})$ should have a component $\mathbf{c}_j^p \in \Delta_j^p = \Gamma_j$, $j \neq i$, contradiction. Therefore $Cl(\mathbf{Q}_i^{\sigma-\mu_i}) \in \Delta_i$. Then, from $\mathbf{Q}_i^{p(\sigma-\mu_i)}$ principal, it follows that $Cl(\mathbf{Q}_i^{\sigma-\mu_i}) \in \Gamma_i$ because Δ_i is cyclic of order p^2 .
- (c) Thus there exists a_i , $1 \leq a_i \leq p-1$, such that $Cl(\mathbf{Q}_i^{\sigma-\mu_i}) = Cl(\mathbf{Q}_i^{a_i p})$ and so $\mathbf{Q}_i^{\sigma-\mu_i-a_i}$ is principal.

In this section we examine the case of subgroups Δ_i of order p^2 . Let us note Δ for one of this groups annihilated by $\sigma - (\mu + ap)$, $a \neq 0$.

Theorem 3.4. μ verifies the two congruences

$$(15) \quad \begin{aligned} \sum_{i=0}^{p-2} \mu^i \delta_i &\equiv 0 \pmod{p} \text{ with } \delta_i = \frac{v_{-(i-1)} - v_{-i}v}{p}, \\ \sum_{i=0}^{p-2} \mu^{p-2+i} \delta_i + (\mu^{p-1} - 1) \sum_{i=1}^{p-2} i \mu^{i-1} \delta_i &\equiv 0 \pmod{p^2}. \end{aligned}$$

Proof.

1. There exists prime ideals $\mathbf{Q}$ of O_K with $Cl(\mathbf{Q}) \in \Delta$, hence $\mathbf{Q}^{p^2}$ principal and $\mathbf{Q}^p$ not principal. From Stickelberger relation $g(\mathbf{Q})^p O_K = \mathbf{Q}^{P(\sigma)}$ where $P(\sigma)$ has been defined in lemma 2.2 p. 4. Then $g(\mathbf{Q})^{p(\sigma-v)} O_K = \mathbf{Q}^{P(\sigma)(\sigma-v)}$, hence from lemma 2.4 p. 4 we get $g(\mathbf{Q})^{p(\sigma-v)} O_K = \mathbf{Q}^{pQ(\sigma)}$, hence $g(\mathbf{Q})^{(\sigma-v)} O_K = \mathbf{Q}^{Q(\sigma)}$. We know, for instance from Ribenboim [9] F. p. 440 that $g(\mathbf{Q})^{\sigma-v} \in K$ so $\mathbf{Q}^{Q(\sigma)}$ is principal. But $\mathbf{Q}^{\sigma-(\mu+ap)}$ is principal hence $\mathbf{Q}^{Q(\mu+ap)}$ is principal, and thus

$$(16) \quad Q(\mu + ap) \equiv 0 \pmod{p^2}.$$

2. From lemma 2.4 p. 4, $Q(\sigma) = \sum_{i=0}^{p-2} \sigma^i \delta_i$ where $\delta_i = \frac{v_{-(i-1)} - v_{-i}v}{p}$. From relation (16)

$$\sum_{i=0}^{p-2} (\mu + ap)^i \delta_i \equiv 0 \pmod{p^2},$$

hence

$$\sum_{i=0}^{p-2} \mu^i \delta_i + ap \sum_{i=1}^{p-1} i \mu^{i-1} \delta_i \equiv 0 \pmod{p^2}.$$

From theorem 3.1 p. 6 applied to the ideal $\mathbf{Q}^p \in \Gamma$ of order p ,

$$\sum_{i=0}^{p-2} \mu^i \delta_i \equiv 0 \pmod{p}.$$

$\mathbf{Q}^{\sigma^{p-1}-1}$ is principal, therefore $\mathbf{Q}^{(\mu+ap)^{p-1}-1}$ is principal and so $(\mu+ap)^{p-1}-1 \equiv 0 \pmod{p^2}$, hence $\mu^{p-1} + (p-1)\mu^{p-2}ap - 1 \equiv 0 \pmod{p^2}$, hence $\mu^{p-1} - 1 - \mu^{p-2}ap \equiv 0 \pmod{p^2}$, hence $a \equiv \frac{\mu^{p-1}-1}{\mu^{p-2}p} \pmod{p}$, and so

$$\sum_{i=0}^{p-2} \mu^i \delta_i + \frac{\mu^{p-1}-1}{\mu^{p-2}} \sum_{i=1}^{p-1} i \mu^{i-1} \delta_i \equiv 0 \pmod{p^2}$$

and finally

$$(17) \quad \sum_{i=0}^{p-2} \mu^{p-2+i} \delta_i + (\mu^{p-1}-1) \sum_{i=1}^{p-2} i \mu^{i-1} \delta_i \equiv 0 \pmod{p^2},$$

which achieves the proof. □

Example:

1. This congruence $\pmod{p^2}$ is valid for no irregular prime numbers $p < 4001$ with rank r of C_p verifying $r > 1$ (verified with a MAPLE program). Therefore the class group of K has no cyclic subgroups of order p^2 for the primes
 $p = 157, 353, 379, 467, 491, 547, 587, 617, 647, 673, 691, 809, 929, 1151, 1217,$
 $1291, 1297, 1307, 1663, 1669, 1733, 1789, 1847, 1933, 1997, 2003, 2087,$
 $2273, 2309, 2371, 2383, 2423, 2441, 2591, 2671, 2789, 2909, 2939, 2957,$
 $3391, 3407, 3511, 3517, 3533, 3539, 3559, 3593, 3617, 3637, 3833, 3851, 3881.$
2. (see table of irregular prime in Washington [11] p. 410). Our result is consistent for the primes $p = 157, 353, 467, 491$ with Schoof [10] table 4.2 p. 1239 describing structure of class groups of some p -cyclotomic fields.

4 On prime factors $h \neq p$ of the class number of the p -cyclotomic field

In previous sections we considered the relative p -class group C_p^- of K . By opposite, in this section we apply Stickelberger relation to all the primes $h \neq p$ dividing the class number $h(K)$. A first subsection is devoted to the general case of the relative class group $\mathbf{C}^-$ of K , a second to the class group of the quadratic subfield of K and a third subsection to the class group to the biquadratic subfield of K when $p \equiv 1 \pmod{4}$.

4.1 The general case

1. The class group $\mathbf{C}$ of K is the direct sum of the class group $\mathbf{C}^+$ of the maximal totally real subfield K^+ of K and of the relative class group $\mathbf{C}^-$ of K .
2. Remind that v is a primitive root mod p and that v_n is to be understood as $v^n \bmod p$ with $1 \leq v_n \leq p-1$. Let $h(K)$ be the class number of K . Let $h \neq p$ be an odd prime dividing $h(K_p)$, with $v_h(h(K)) = \beta$. Let $d = \text{Gcd}(h-1, p-1)$. Let $\mathbf{C}(h)$ be the h -Sylow subgroup of the class group of K of order h^β . Then $\mathbf{C}(h) = \bigoplus_{j=1}^{\rho} \mathbf{C}_j(h)$ where ρ is the h -rank of the abelian group $\mathbf{C}(h)$ of order h^β and $\mathbf{C}_j(h)$ are cyclic groups of order h^{β_j} where $\beta = \sum_{i=1}^{\rho} \beta_j$.
3. From Kummer (see for instance Ribenboim [8] (3A) p. 119), the prime ideals of O_K of inertial degree 1 generate the ideal class group. Therefore there exist in the subgroup C_h of exponent h of $\mathbf{C}(h)$ some prime ideals $\mathbf{q}$ of inertial degree 1 such that $Cl(\mathbf{q}) \in \bigoplus_{j=1}^{\rho} c_j$ where c_j is a cyclic group of order h and $Cl(\mathbf{q}) \notin \bigoplus_{j \in J} c_j$ where J is a strict subset of $\{1, 2, \dots, \rho\}$.
4. Let $P(\sigma) = \sum_{k=0}^{p-2} \sigma^k v_{-k}$ be the Stickelberger polynomial. From lemma 2.2 p. 4 Stickelberger relation is $\mathbf{q}^{P(\sigma)} = g(\mathbf{q})^p O_K$ where $g(\mathbf{q})^p \in O_K$. Therefore $\mathbf{q}^{P(\sigma)}$ is principal, a fortiori is $\mathbf{C}(h)$ -principal (or $Cl(\mathbf{q})^{P(\sigma)}$ has a trivial component in $\mathbf{C}(h)$). There exists a minimal polynomial $V(X) \in \mathbf{F}_h(X)$ of degree $\delta \leq \rho$ such that $\mathbf{q}^{V(\sigma)}$ is $\mathbf{C}(h)$ -principal, if not the remainder of the division of $P(X)$ by $V(X)$ of degree smaller than δ would annihilate also $\mathbf{q}$. Therefore the irreducible polynomial $V(X)$ divides $P(X)$ in $\mathbf{F}_h[X]$ for the indeterminate X .
5. If $Cl(\mathbf{q}) \in \mathbf{C}^-$ then $\mathbf{q}^{\sigma^{(p-1)/2}+1}$ is principal.
6. Let $D(X) \in \mathbf{F}_h(X)$ defined by $D(X) = \text{Gcd}(P(X), X^{(p-1)/2} + 1) \bmod h$.

We obtain the following:

Theorem 4.1. *Suppose that the prime $h \neq p$ divides the class number $h(K)$. Let $D(X) = \text{Gcd}(P(X), X^{(p-1)/2} + 1) \bmod h$. Then*

1. $V(X)$ divides $D(X)$ in $\mathbf{F}_h[X]$. The h -rank ρ of $\mathbf{C}(h)$ is greater or equal to the degree δ of $V(X)$ and $h^\rho \mid h(K)$.
2. If h is coprime with $p-1$ and with the class number $h(E)$ of all intermediate fields $\mathbb{Q} \subset E \subset K$, $E \neq K$ then $f \mid \rho$ where f is the order of $h \bmod p-1$.

Proof.

1. Reformulation of previous paragraph.
2. Immediate consequence of theorem 10.8 p. 187 in Washington [11] for the cyclic extension K/E .

□

Remark: See the section 5 p. 18 for a MAPLE program applying the theorem 4.1 p. 12.

Corollary 4.2. *If $\mathbf{C}(h)$ is cyclic then:*

1. $V(X) = X - \nu$ with $\nu \in \mathbf{F}_h^*$.
2. In $\mathbf{F}_h(X)$

$$(18) \quad \begin{aligned} & V(X) \mid X^d - 1, \quad d = \gcd(h-1, p-1), \\ & V(X) \mid \left\{ \sum_{i=0}^{d-1} X^i \times \frac{\sum_{j=0}^{(p-1)/d-1} v^{-(i+jd)}}{p} \right\}. \end{aligned}$$

3. Let M be the smallest subfield of K such that $h \mid h(M)$. Let $n = [M : \mathbb{Q}]$. If h is coprime with n then $h-1 \equiv 0 \pmod{n}$.

Proof.

1. $\rho = 1$ implies that $V(\sigma) = \sigma - \nu$.
2. $X - \nu \mid X^{h-1} - \nu^{h-1}$ and so $\sigma^{h-1} - \nu^{h-1}$ annihilates $\mathbf{C}(h)$. From $\nu^{h-1} \equiv 1 \pmod{h}$ it follows that $\sigma^{h-1} - 1$ annihilates $\mathbf{C}(h)$. $\sigma^{p-1} - 1$ annihilates $\mathbf{C}(h)$ and so $\sigma^d - 1$ annihilates $\mathbf{C}(h)$ and so $X - \nu$ divides $X^d - 1$ in $\mathbf{F}_h[X]$. Then apply theorem 4.1. Observe that $\frac{\sum_{j=0}^{(p-1)/d-1} v^{-(i+jd)}}{p} \equiv 0 \pmod{p}$ and that we have assumed that the prime $h \neq p$ in this section.
3. Let E be the smallest intermediate field E , $\mathbb{Q} \subset E \subset K$ with $h \mid h(E)$ and $[E : \mathbb{Q}] = n$. Then we apply theorem 10.8 p. 187 of Washington [11] to the cyclic extension $M/\mathbb{Q}$, thus $h \equiv 1 \pmod{n}$ and $n \mid p-1$ because $f \mid \rho$ where $\rho = 1$.

□

It is possible to enlarge previous results with another annihilation polynomial:

Theorem 4.3. *The polynomial $\Pi(\sigma) = \sum_{i=0, v_{-i} \text{ even}}^{p-2} \sigma^i$ annihilates the non- p -part of the class group $\mathbf{C}$ of the p -cyclotomic field K .*

Proof.

1. Apply Stickelberger relation to field $\mathbb{Q}(\zeta_{2p}) = \mathbb{Q}(\zeta_p)$. Let $\varpi_{2t+1} : \zeta_{2p} \rightarrow \zeta_{2p}^{2t+1}$. The Stickelberger polynomial can be written

$$S_2 = \sum_{2t+1=1, t \neq p}^{2p-1} \varpi_{2t+1}^{-1} \times (2t+1).$$

2. Observe at first that $\zeta_{2p} = -\zeta_p$.

If $t > (p-1)/2$

then $\varpi_{2t+1} : \zeta_{2p} \rightarrow -\zeta_{2p}^{2t+1-p}$, hence $\varpi_{2t+1} : -\zeta_p \rightarrow -(-\zeta_p)^{2t+1-p}$, hence $\varpi_{2t+1} : -\zeta_p \rightarrow -(\zeta_p)^{2t+1-p}$ because $2t+1-p$ is even, hence $\varpi_{2t+1} : \zeta_p \rightarrow \zeta_p^{2t+1-p}$, hence $\varpi_{2t+1} = \varpi_{2t+1-p}$, hence

$$\varpi_{2t+1}^{-1} \times (2t+1) = \varpi_{2t+1-p}^{-1} \times (2t+1-p) + p \times \varpi_{2t+1-p}^{-1}.$$

3. The Stickelberger polynomial is

$$S_2 = \sum_{t=0}^{(p-3)/2} \varpi_{2t+1}^{-1} \times (2t+1) + \sum_{t=(p+1)/2}^{p-1} \varpi_{2t+1}^{-1} \times (2t+1),$$

hence

$$S_2 = \sum_{t=0}^{(p-3)/2} \varpi_{2t+1}^{-1} \times (2t+1) + \sum_{t=(p+1)/2}^{p-1} \varpi_{2t+1-p}^{-1} \times (2t+1-p) + p \sum_{t=(p-1)/2}^{p-1} \varpi_{2t+1-p},$$

hence

$$S_2 = \sum_{t=1}^{p-1} \varpi_t^{-1} t + p \times \sum_{t=(p-1)/2}^{p-1} \varpi_{2t+1-p},$$

hence, with $P(\sigma)$ defined in lemma 2.2 p. 4,

$$S_2 = P(\sigma) + p \times \sum_{t=(p-1)/2}^{p-1} \varpi_{2t+1-p},$$

With $2t+1-p = v_{-i}$ we get

$$(19) \quad S_2 = P(\sigma) + p \times \sum_{i=0, v_{-i} \text{ even}}^{p-2} \sigma^i.$$

4. The polynomial $P(\sigma)$ annihilates the class group $\mathbf{C}$ of K . Therefore the polynomial $p \times \sum_{i=0, v_{-i} \text{ even}}^{p-2} \sigma^i$ annihilates also $\mathbf{C}$. If $h \neq p$ then $\Pi(\sigma)$ annihilates $\mathbf{C}(h)$, which achieves the proof.

□

Remark: Numerical MAPLE computations seem to show more : the polynomial

$$\text{Gcd}(\Pi(\sigma), \sigma^{(p-1)/2} + 1) = \frac{\sigma^{(p-1)/2} + 1}{\sigma - v} = \prod_{m=1}^{(p-3)/2} (\sigma - v_{2m+1}).$$

Therefore $\Pi(\sigma)$ annihilates also the relative p -class group C_p^- .

Lemma 4.4. *Let E be a subfield of K with $[K : \mathbb{Q}] = d$. Let h be an odd prime number dividing $h(L)$. Then in $\mathbf{F}_h[X]$*

$$(20) \quad \begin{aligned} V(X) &| \sum_{i=0}^{d-1} X^i, \\ V(X) &| \left\{ \sum_{i=0}^{d-1} X^i \times \frac{\sum_{j=0}^{(p-1)/d-1} v^{-(i+jd)}}{p} \right\}. \end{aligned}$$

Proof. $\sigma^d - 1$ annihilates $\mathbf{C}(h)$. The Stickelberger polynomial

$$P(X) = \sum_{i=0}^{p-2} X^i v_{-i} = p \times \sum_{i=0}^{d-1} X^i \times \frac{\sum_{j=0}^{(p-1)/d-1} v^{-(i+jd)}}{p},$$

and from $p \neq h$ it follows that

$$V(X) | \left\{ \sum_{i=0}^{d-1} X^i \times \frac{\sum_{k=0}^{(p-1)/d-1} v^{-(i+jd)}}{p} \right\}$$

in $\mathbf{F}_p[G]$. □

Remark: Compare lemma 4.4 for $h \neq p$ with lemma 3.2 p. 7 proved when $h = p$.

4.2 The case of complex quadratic fields contained in K

In this paragraph we formulate directly previous result when h divides the class number of the complex quadratic field $\mathbb{Q}(\sqrt{-p}) \subset K$, $p \equiv 3 \pmod{4}$, $p \neq 3$.

Theorem 4.5. *Hilbert 145 theorem*

Suppose that $p \equiv 3 \pmod{4}$, $p \neq 3$. If h is an odd prime with $h \mid h(\mathbb{Q}(\sqrt{-p}))$ then

$$(21) \quad \sum_{i=0}^{p-2} (-1)^i v_{-i} \equiv 0 \pmod{h}.$$

Proof. Let $\mathbf{Q}$ be the prime of $\mathbb{Q}(\sqrt{-p})$ lying above $\mathbf{q}$. The ideals $\mathbf{Q} \neq \sigma(\mathbf{Q})$ and so $\mathbf{Q}^{\sigma+1}$ is principal because $\mathbf{Q}^{\sigma^2} = \mathbf{Q}$. Therefore $\mathbf{Q}^{\sum_{i=0}^{p-2} (-1)^i v_{-i}}$ is principal and $\sum_{i=0}^{p-2} (-1)^i v_{-i} \equiv 0 \pmod{h}$. □

Remarks:

1. The theorem 4.7 can also be obtained from Hilbert Theorem 145 see Hilbert [3] p. 119. See also Mollin, [6] theorem 5.119 p. 318.
2. From lemmas 2.4 p. 4 we could prove similarly:
Suppose that $p \equiv 3 \pmod{4}$, $p \neq 3$. If h is an odd prime with $h \mid h(\mathbb{Q}(\sqrt{-p}))$ then

$$(22) \quad 2 \times \sum_{i=0}^{(p-3)/2} (-1)^i v_{-i} - p \equiv 0 \pmod{h}.$$

3. Numerical evidences easily computable show more: If $p \neq 3$ is prime with $p \equiv 3 \pmod{4}$ then the class number $h(\mathbb{Q}(\sqrt{-p}))$ verifies

$$(23) \quad h(\mathbb{Q}(\sqrt{-p})) = -\frac{\sum_{i=0}^{p-2} (-1)^i v_{-i}}{p}.$$

This result has been proved by Dirichlet by analytical number theory, see Mollin remark 5.124 p. 321. It is easy to verify this formula, for instance in tables of class numbers of complex quadratic fields in :

- (a) H. Cohen [2] p. 502- 505, all the table for $p \leq 503$.
- (b) in Wolfram table of quadratic class numbers [12] for large p .
4. When $p \equiv 1 \pmod{4}$ this method cannot be applied to the quadratic field $\mathbb{Q}(\sqrt{p}) \subset K$ because $\sum_{i=0}^{p-2} (-1)^i v_{-i}$ is trivially null.

Theorem 4.6. *Suppose that $p \equiv 3 \pmod{4}$, $p \neq 3$. If h is an odd prime with $h \mid h(\mathbb{Q}(\sqrt{-p}))$ then*

$$(24) \quad \begin{aligned} & \sum_{i=0, v_{-i} \text{ even}}^{p-2} (-1)^i \neq 0, \\ & \sum_{i=0, v_{-i} \text{ even}}^{p-2} (-1)^i \equiv 0 \pmod{h}. \end{aligned}$$

Proof. We apply previous theorem 4.3 p. 13 observing that in that case $\sigma^2(\mathbf{Q}) = \mathbf{Q}$ when $\mathbf{Q}$ is a non-principal ideal of the quadratic field $\mathbb{Q}(\sqrt{-p})$, hence $\sigma + 1$ annihilates $\mathbf{C}(h)$ and $\sum_{i=0, v_{-i} \text{ even}}^{p-2} (-1)^i \equiv 0 \pmod{h}$. This sum has $\frac{p-1}{2}$ elements, thus of odd cardinal and cannot be null. $\square$

Theorem 4.7. Suppose that $p \equiv 3 \pmod{4}$, $p \neq 3$. Let δ be an integer $1 \leq \delta \leq p-2$. Let I_δ be the set

$$(25) \quad I_\delta = \{i \mid 0 \leq i \leq p-2, v_{(p-1)/2-i} + v_{(p-1)/2-i+ind_v(\delta)} > p\},$$

where, as seen above, $ind_v(\delta)$ is the notation index of δ relative to v . If h is an odd prime with $h \mid h(\mathbb{Q}(\sqrt{-p}))$ then

$$(26) \quad \begin{aligned} \sum_{i \in I_\delta} (-1)^i &\neq 0, \\ \sum_{i \in I_\delta} (-1)^i &\equiv 0 \pmod{h}. \end{aligned}$$

Proof. I_δ has an odd cardinal. Then see relation (25). $\square$

Remark:

1. Observe that results of theorems 4.6 and 4.7 are consistent with existing tables of quadratic fields, for instance Arno, Robinson, Wheeler [1]. Numerical verifications seem to show more :

$$(27) \quad \sum_{i=0, v_{-i} \text{ even}}^{p-2} (-1)^i \equiv 0 \pmod{h(\mathbb{Q}(\sqrt{-p}))}.$$

2. Observe that if $p \equiv 1 \pmod{4}$ then $\sum_{i=0, v_{-i} \text{ even}}^{p-2} (-1)^i = 0$.

4.3 The case of biquadratic fields contained in K

The following example is a generalization for the biquadratic fields L which are included in p -cyclotomic field K with $p \equiv 1 \pmod{4}$.

Theorem 4.8. Let p be a prime with $2^2 \parallel p-1$. Let

$$(28) \quad S = \left(\sum_{i=0}^{(p-3)/2} (-1)^i v_{2i} \right)^2 + \left(\sum_{i=0}^{(p-3)/2} (-1)^i v_{2i+1} \right)^2.$$

Let L be the field with $\mathbb{Q}(\sqrt{p}) \subset L \subset K$, $[L : \mathbb{Q}(\sqrt{p})] = 2$. Let h be an odd prime number with $h \mid h(L)$ and $h \nmid h(\mathbb{Q}(\sqrt{p}))$. Then $S \neq 0$ and $S \equiv 0 \pmod{h}$.

Proof. $V(\sigma) \mid \sigma^4 - 1$. $h \nmid h(\mathbb{Q}(\sqrt{p}))$ and so $V(\sigma) \mid \sigma^2 + 1$. $P(\sigma) = \sum_{i=0}^{(p-3)/2} \sigma^{2i} v_{-2i} + \sigma \times \sum_{i=0}^{(p-3)/2} \sigma^{2i} v_{2i+1}$. Relation (28) follows. $\square$

Remarks:

1. S does not depend of the primitive root $v \bmod p$ chosen.
2. Numerical computations seem to show more : $P(\sigma) \equiv 0 \bmod p^2$ and so

$$(29) \quad \frac{(\sum_{i=0}^{(p-3)/2} (-1)^i v_{2i})^2 + (\sum_{i=0}^{(p-3)/2} (-1)^i v_{2i+1})^2}{p^2} \equiv 0 \bmod h.$$

3. This result is a generalization for biquadratic fields of theorem 145 of Hilbert for quadratic fields.

5 A numerical MAPLE algorithm

This section contains a MAPLE algorithm connected to the structure of the relative class group $\mathbf{C}^-$. For each prime number $p < 500$, the algorithm computes

1. a primitive root $v \bmod p$,
2. the Stickelberger polynomial $P(X)$ has been defined in lemma 2.2 p. 4 and $Q(X)$ in lemma 2.4 p. 4. For the prime numbers h with $3 \leq h \leq p^2$ and the primitive root $v \bmod p$, the algorithm computes the polynomial $GCD(X) \in \mathbf{F}_h[X]$ for the indeterminate X given by formulas:

(a) if $h \neq p$ then

(b)

$$(30) \quad GCD(X) = Gcd(P(X), X^{(p-1)/2} + 1) \bmod h,$$

(c) if $h = p$ then

$$(31) \quad GCD(X) = Gcd(Q(X), X^{(p-1)/2} + 1) \bmod h,$$

The results are compared with the corresponding tables of class numbers of cyclotomic fields $K = Q(\zeta_p)$ for the primes $p < 500$ in Schoof [10]. We observe that:

1. The set of odd prime numbers h with $\deg(GCD(X)) > 0$ is strictly the set of odd prime divisors h of K in Schoof table p. 1142.
2. The rank ρ of the h -Sylow subgroup $\mathbf{C}(h)$ in Schoof tables 4.2 p. 1239 (h not dividing $p-1$) and 4.3 p. 1240 (h dividing $p-1$) is the degree of the polynomial $GCD(X)$ found here. Observe that when $h(K^+) \not\equiv 0 \bmod p$ and $h = p$ this fact can be proved by means out of reach of this article at elementary level (Ribet theorem).

3. Let $GCD(X) = \prod_{i=1}^n A_i(X)^{n_i}$ be the prime decomposition of $GCD(X)$ in the euclidean field $\mathbf{F}_h[X]$. We observe that to each prime polynomial $A_i(X)$ corresponds a subgroup of $\mathbf{C}(h)$ of h -rank $d_i \times n_i$ where d_i is the degree of $A_i(X)$. In particular if $d_i = n_i = 1$ then the subgroup corresponding to $A_i(X)$ is cyclic. Note that for all prime p with $h(K^+) \not\equiv 0 \pmod{p}$ and $h = p$ it is a consequence of Ribet theorem.
4. We observe that when $h \neq p$ there exists some cases where $n_i > 1$ for instance for $p = 101$ and $h = 5$ with $n_1 = 2$. In these cases this implies that the minimal polynomial $V(X)$ annihilating C_h is different of $GCD(X) : V(X) \mid GCD(X)$ and $\text{degree}(V(X)) < \text{degree}(GCD(X))$.

A question: for all the odd primes $p < 500$ and all the odd primes $h < p^2$ we have observed that the degree of $GCD(X)$ is equal to the rank ρ of the h -Sylow subgroup $\mathbf{C}(h)$ of the relative class group $\mathbf{C}^-$ of K with $GCD(X) = 1 \Leftrightarrow h(K) \not\equiv 0 \pmod{h}$: this gives important informations on the structure of the relative class group $\mathbf{C}^-$ of K : the precise set of odd primes h dividing $h(K)$ and for each of them the rank ρ of the h -group $\mathbf{C}(h)$. Can we generalize this property to all the odd primes p and all the odd primes h or at least at some predefined subsets of them?

The MAPLE algorithm

```

restart;
> p:=3:
> while p<499 do
> p:=nextprime(p):
> for v from 2 to p-2 do:
> i_v:=1:
> for i from 2 to p-2 do:
> if v&^i mod p = 1 then i_v:=0: fi:
> od:
> if i_v=1 then
> T:= X^((p-1)/2)+1 :
> S:=0:
> Q:=0:
> for i from 0 to p-2 do:
> vmi:=v&^(p-1-i) mod p:
> vmim1:=v&^(p-1-(i-1)) mod p:
> delta_i:=iquo(vmim1-v*vmi,p):
> S:=S+X^i*vmi:
> Q:=Q+X^i*delta_i:
> od:
> for h from 3 to p^2 do:
> if isprime(h)=true then
> if h<>p then
> GCD:=Gcd(T,S) mod h:
> deg_GCD:= degree (GCD):
> if deg_GCD>0 then
> GCD_Factors:=Factors(GCD) mod h:

```

```

> n:=nops(GCD_Factors[2]):
> rho:=0:
> for i from 1 to n do:
> temp:=GCD_Factors[2]:
> temp:=temp[i]:
> temp1:=temp[1]:
> temp2:=temp[2]:
> rho:=rho+degree(temp1)*temp2:
> od:
> print('p=',p,'h=',h,'rho=',rho,'v=',v,'GCD(X)=' ,GCD_Factors[2]):
> fi:
> fi:
> if h = p then
> GCD1:=Gcd(T,Q) mod h:
> deg_GCD1:= degree (GCD1):
> if deg_GCD1>0 then
> GCD1_Factors:=Factors(GCD1) mod h:
> n:=nops(GCD1_Factors[2]):
> rho_1:=0:
> for i from 1 to n do:
> temp:=GCD1_Factors[2]:
> temp:=temp[i]:
> temp1:=temp[1]:
> temp2:=temp[2]:
> rho_1:=rho_1+degree(temp1)*temp2:
> od:
> print('p=',p,'h=',h,'rho_1=',rho_1,'v=',v,'GCD1=' ,GCD1_Factors[2]):
> fi:
> fi:
> fi:
> od:
> break:
> fi:
> od:
> #fi:
> od:
>

```

In the following page the table of results obtained.

p	h	ρ	v	$GCD(X)$
23	3	1	5	$X + 1$
31	3	1	3	$X + 1$
37	37	1	2	$X + 5$
41	11	2	6	$X^2 + 10X + 6$
43	211	1	3	$X + 73$
47	5	1	5	$X + 1$
47	139	1	5	$X + 44$
59	3	1	2	$X + 1$
59	59	1	2	$X + 36$
59	233	1	2	$X + 8$
61	41	1	2	$X + 36$
61	1861	1	2	$X + 997$
67	67	1	2	$X + 24$
71	7	1	7	$X + 1$
73	89	1	5	$X + 12$
79	5	1	3	$X + 1$
79	53	1	3	$X + 28$
83	3	1	2	$X + 1$
89	113	1	3	$X + 95$
97	577	1	5	$X + 46$
97	3457	1	5	$X + 1558$
101	5	2	2	$(X + 3)^2$
101	101	1	2	$X + 66$
101	601	1	2	$X + 323$
103	5	1	5	$X + 1$
103	103	1	5	$X + 58$
103	1021	1	5	$X + 9$
107	3	1	2	$X + 1$
107	743	1	2	$X + 50$
107	9859	1	2	$X + 4936$
109	17	1	6	$X + 4$
109	1009	1	6	$X + 41$
113	17	1	3	$X + 5$
127	5	1	3	$X + 1$
127	13	1	3	$X + 9$
127	43	1	3	$X + 4$
127	547	1	3	$X + 169$
127	883	1	3	$X + 336$
127	3079	1	3	$X + 1925$
131	3	3	2	$X^3 + 2X^2 + 1$
131	5	1	2	$X + 1$
131	53	1	2	$X + 46$
131	131	1	2	$X + 34$
131	1301	1	2	$X + 283$
137	17	1	3	$X + 8$
139	3	1	2	$X + 1$
139	47	1	2	$X + 9$
139	277	2	2	$(X + 191)(X + 218)$
139	967	1	2	$X + 241$
149	3	2	2	$X^2 + 1$
149	149	1	2	$X + 43$
151	7	1	6	$X + 1$
151	11	2	6	$X^2 + 6X + 3$
151	281	1	6	$X + 90$

p	h	ρ	v	$GCD(X)$
157	5	1	5	$X + 3$
157	13	1	5	$X + 6$
157	157	2	5	$(X + 95)(X + 91)$
157	1093	1	5	$X + 800$
157	1873	1	5	$X + 935$
163	181	1	2	$X + 65$
163	23167	1	2	$X + 8783$
167	11	1	5	$X + 1$
167	499	1	5	$X + 491$
173	5	1	2	$X + 2$
173	20297	1	2	$X + 997$
179	5	1	2	$X + 1$
179	1069	1	2	$X + 552$
181	5	1	2	$X + 3$
181	37	1	2	$X + 29$
181	41	1	2	$X + 2$
181	61	1	2	$X + 6$
181	1321	1	2	$X + 149$
181	2521	1	2	$X + 2015$
191	11	1	19	$X + 3$
191	13	1	19	$X + 1$
193	6529	1	5	$X + 4193$
193	15361	1	5	$X + 13057$
193	29761	1	5	$X + 29163$
197	5	1	2	$X + 3$
197	1877	1	2	$X + 981$
197	7841	1	2	$X + 1604$
199	3	1	3	$X + 1$
199	19	1	3	$X + 4$
199	727	1	3	$X + 590$
211	3	1	2	$X + 1$
211	7	1	2	$X + 4$
211	41	1	2	$X + 16$
211	71	1	2	$X + 15$
211	181	1	2	$X + 5$
211	281	2	2	$(X + 109)(X + 101)$
211	421	1	2	$X + 93$
211	1051	1	2	$X + 884$
211	12251	1	2	$X + 1580$
223	7	1	3	$X + 1$
223	43	1	3	$X + 36$
227	5	1	2	$X + 1$
227	2939	3	2	$(X + 1420)(X + 509)(X + 2006)$
229	13	1	6	$X + 6$
229	17	1	6	$X + 13$
229	457	1	6	$X + 126$
229	7753	1	6	$X + 4310$
233	233	1	3	$X + 193$
233	1433	1	3	$X + 1091$
239	3	1	7	$X + 1$
239	5	1	7	$X + 1$

p	h	ρ	v	$GCD(X)$	p	h	ρ	v	$GCD(X)$
241	47	2	7	$X^2 + 29X + 1$	353	353	2	3	$(X + 299)(X + 51)$
241	13921	1	7	$X + 9953$	353	6113	1	3	$X + 2060$
241	15601	1	7	$X + 7049$	353	9473	1	3	$X + 5067$
251	7	1	6	$X + 1$	359	19	1	7	$X + 1$
251	11	1	6	$X + 4$	367	3	1	6	$X + 1$
257	257	1	3	$X + 76$	367	733	1	6	$X + 686$
263	13	1	5	$X + 1$	367	39163	1	6	$X + 27454$
263	263	1	5	$X + 204$	373	5	1	2	$X + 2$
263	787	1	5	$X + 510$	373	61	1	2	$X + 21$
269	13	1	2	$X + 5$	373	1117	1	2	$X + 532$
271	11	1	6	$X + 1$	373	1489	1	2	$X + 990$
271	31	1	6	$X + 2$	379	3	1	2	$X + 1$
271	37	1	6	$X + 9$	379	13	1	2	$X + 3$
271	271	1	6	$X + 196$	379	127	1	2	$X + 13$
271	811	1	6	$X + 133$	379	379	2	2	$(X + 348)(X + 91)$
271	1201	1	6	$X + 367$	379	547	1	2	$X + 196$
271	1621	1	6	$X + 1190$	379	757	1	2	$X + 531$
271	15391	1	6	$X + 6331$	379	991	1	2	$X + 324$
271	21961	1	6	$X + 6698$	379	1499	1	2	$X + 1314$
277	17	1	5	$X + 13$	379	9199	1	2	$X + 7901$
277	47	2	5	$X^2 + 19X + 42$	383	17	1	5	$X + 1$
277	829	1	5	$X + 150$	389	41	1	2	$X + 32$
281	11	2	3	$X^2 + 2X + 2$	389	389	1	2	$X + 231$
281	17	1	3	$X + 8$	389	1553	1	2	$X + 1130$
281	41	1	3	$X + 24$	397	13	1	5	$X + 5$
281	401	1	3	$X + 250$	397	23	2	5	$X^2 + 13X + 18$
283	3	1	3	$X + 1$	397	109	1	5	$X + 32$
283	283	1	3	$X + 236$	397	4861	1	5	$X + 3655$
293	3	2	2	$X^2 + 1$	397	9901	1	5	$X + 8544$
293	293	1	2	$X + 194$	401	41	2	3	$X^2 + 17$
307	3	1	5	$X + 1$	401	401	1	3	$X + 141$
307	37	1	5	$X + 33$	401	64849	1	3	$X + 46775$
307	137	1	5	$X + 38$	409	5	2	21	$X^2 + 3$
307	307	1	5	$X + 16$	409	17	1	21	$X + 2$
307	443	1	5	$X + 13$	409	73	1	21	$X + 66$
307	613	1	5	$X + 49$	409	409	1	21	$X + 28$
307	919	1	5	$X + 144$	409	1321	1	21	$X + 1304$
307	1429	1	5	$X + 1294$	419	3	1	2	$X + 1$
311	19	1	17	$X + 1$	419	1103	1	2	$X + 494$
311	41	1	17	$X + 18$	421	5	1	2	$X + 2$
311	311	1	17	$X + 158$	421	29	1	2	$X + 18$
313	37	2	10	$X^2 + 14$	421	37	1	2	$X + 23$
313	233	1	10	$X + 136$	421	421	1	2	$X + 72$
317	13	1	2	$X + 5$	421	2521	1	2	$X + 60$
331	3	6	3	$(X + 1)^2(X^4 + 2X^3 + X^2 + 2X + 1)$	421	39509	1	2	$X + 7582$
331	23	1	3	$X + 2$	421	39901	1	2	$X + 8081$
331	61	1	3	$X + 12$	421	70309	1	2	$X + 65038$
331	67	1	3	$X + 64$	431	3	1	7	$X + 1$
337	7	2	10	$X^2 + X + 6$	431	7	1	7	$X + 1$
337	17	2	10	$(X + 10)(X + 7)$	431	11	1	7	$X + 5$
337	353	1	10	$X + 36$	431	701	1	7	$X + 210$
347	5	1	2	$X + 1$	431	14621	1	7	$X + 2522$
347	347	1	2	$X + 52$	433	433	1	5	$X + 371$
349	5	1	2	$X + 2$	433	3457	1	5	$X + 2700$
349	13	1	2	$X + 6$	433	12097	1	5	$X + 31$
349	2089	1	2	$X + 1733$	433	21601	1	5	$X + 10658$
349	17749	1	2	$X + 9289$	433	47521	1	5	$X + 36247$

p	h	ρ	v	$GCD(X)$
439	3	2	15	$(X+1)^2$
439	5	1	15	$X+1$
439	293	1	15	$X+283$
443	3	3	2	X^3+X^2+2X+1
443	5	1	2	$X+1$
443	79	1	2	$X+8$
443	157	1	2	$X+67$
443	12377	1	2	$X+6026$
449	168449	1	3	$X+33570$
457	5	2	13	X^2+4X+2
457	41	1	13	$X+14$
457	577	1	13	$X+9$
457	1217	1	13	$X+692$
457	43777	1	13	$X+37577$
457	63841	1	13	$X+2827$
461	5	2	2	$(X+2)^2$
461	461	1	2	$X+13$
461	661	1	2	$X+258$
461	161461	1	2	$X+134936$
463	7	2	3	$(X+2)(X+1)$
463	29	1	3	$X+20$
463	89	1	3	$X+64$
463	463	1	3	$X+8$
463	631	1	3	$X+62$
463	673	1	3	$X+223$
463	1123	1	3	$X+49$
463	4423	1	3	$X+387$
463	8779	1	3	$X+5520$
467	7	1	2	$X+1$
467	467	2	2	$(X+239)(X+236)$
479	5	1	13	$X+1$
479	48757	1	13	$X+34844$
479	62141	1	13	$X+43049$
487	7	2	3	$(X+4)(X+1)$
487	37	2	3	$(X+33)(X+12)$
487	919	1	3	$X+267$
487	2647	1	3	$X+1070$
487	10909	1	3	$X+3031$
487	58321	1	3	$X+58241$
491	3	1	2	$X+1$
491	11	2	2	$(X+4)(X+5)$
491	29	1	2	$X+16$
491	491	3	2	$(X+203)(X+419)(X+418)$
499	3	1	7	$X+1$
499	167	1	7	$X+98$

References

- [1] S. Arno, M.L. Robinson, F.S. Wheeler *Imaginary quadratic fields with small odd class number*, Acta Arith. 83, 1998, 295-300.
- [2] H. Cohen, A course in computational number theory, Springer-Verlag, 1993.
- [3] D. Hilbert, *The Theory of Algebraic Numbers*, Springer, 1998

- [4] K. Ireland, M. Rosen, *A Classical Introduction to Modern Number Theory*, Springer-Verlag, 1982.
- [5] H. Koch, *Algebraic Number Theory*, Springer, 1997.
- [6] R.A. Mollin, *Algebraic Number Theory*, Chapman and Hall/CRC, 1999.
- [7] W. Narkiewicz, *Elementary and Analytic Theory of Numbers*, Springer-verlag, 1990.
- [8] P. Ribenboim, *13 Lectures on Fermat's Last Theorem*, Springer-Verlag, 1979.
- [9] P. Ribenboim, *Classical Theory of Algebraic Numbers*, Springer, 2001.
- [10] R. Schoof, *Minus class groups of the fields of the l -roots of unity*, Mathematics of Computation, vol 267,223, jul 1998, p. 1225-1245.
- [11] L.C. Washington, *Introduction to cyclotomic fields, second edition*, Springer, 1997.
- [12] *Class number of quadratic fields*,
<http://www.mathworld.wolfram.com/ClassNumber.html>

Roland Quême

13 avenue du château d'eau
 31490 Brax
 France

mailto:roland.queme@wanadoo.fr

home page: <http://roland.queme.free.fr/>

V10 - MSC Classification : 11R18; 11R29
